



QUADRA

Section 18 Emergency Exemption

SECTION 18 SPECIFIC EMERGENCY EXEMPTION FOR THE USE OF QUADRIS®
FLOWABLE FUNGICIDE ON TOBACCO IN KENTUCKY FROM June 24, 2005 THROUGH
October 15, 2005

File Symbol 05-KY-13

Quadris® Flowable Fungicide

For Control of Certain Diseases in Tobacco

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Follow all application directions, restrictions, and precautions, including statements pertaining to the Worker Protection Standards, on the EPA-registered Quadris Flowable Fungicide label (EPA Reg. No.: 100-1098).

These Use Directions and the EPA-registered Quadris Flowable Fungicide label must be in the possession of the user at the time of pesticide mixing and application.

Integrated Pest (Disease) Management: QUADRIS Flowable Fungicide should be integrated into an overall disease management strategy that includes proper selection of varieties with disease tolerance, removal of plant debris in which inoculum over winters, plant residue management, crop rotation and proper timing and placement of irrigation.

Crop	Target Diseases	Use Rate fl.oz. product/A (lbs. a.i./A)	Remarks
Tobacco	<p>Frog-eye leafspot (<i>Cercospora nicotianae</i>)</p> <p>Target spot (<i>Rhizoctonia solani</i>)</p>	8 (0.13)	<p>Application Directions: QUADRIS Flowable Fungicide applications should begin at the first sign of disease development. Apply on a 7-14 day interval with shorter intervals under conditions conducive to disease development. Do not make more than two consecutive applications of QUADRIS Flowable Fungicide or any other strobilurin fungicide before alternating to a fungicide with a different mode of action for two applications. For ground applications, apply QUADRIS Flowable Fungicide in sufficient water volume for adequate coverage and canopy penetration. For aerial application, volumes should be 10-15 GPA. May be applied up to the day of harvest but should not have more than one application within last 30 days prior to harvest.</p> <p>Resistance Management: QUADRIS Flowable Fungicide must be used in alternation with another fungicide registered on tobacco that controls the target disease(s) including Acrobat MZ or Acrobat + Dithane DF tank mix. Up to four (4) applications of QUADRIS Flowable Fungicide or other strobilurin fungicides may be made per season.</p>

Specific Use Restrictions:

Do not tank mix with Thiodan or any sucker control products.

Do not apply more than 1.1 quart (36 fl. oz/A or 0.60 lb ai/A) per acre per season.

Precautions: QUADRIS Flowable Fungicide may enhance weather flecking on the leaves of certain tobacco types and the grower accepts any liability associated with this risk.

Tank mixing Quadris Flowable Fungicide with insecticides formulated as ECs or containing high amounts of solvents, may cause some crop injury.

Any adverse effects resulting from the use of Quadris Flowable Fungicide under this emergency exemption must be immediately reported to the Kentucky Department of Agriculture.

Quadris is a registered trademark of Syngenta Crop Protection

Syngenta Crop Protection, Inc.
Greensboro, North Carolina 27419

Label Code: KY1098013AB0605